

ICU Pharmacist References

Special Guest: Chris Droege, PharmD, BCCCP, FCCM, FASHP

General/Background Readings

Wijdicks EFM, et al. The early days of the neurosciences intensive care unit. *Mayo Clin Proc* 2011; 86(9): 903-906. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3258005/>

The Ninth Floor Project - <https://pharm.ucsf.edu/history-cp/1965-1972>

Rudis MI, et al. Position paper on critical care pharmacy services. *Crit Care Med* 2000; 28: 3746-3750. <https://www.ncbi.nlm.nih.gov/pubmed/11098984>

Bond CA, et al. Clinical pharmacy services, pharmacy staffing, and hospital mortality rates. *Pharmacotherapy* 2007; 27(4): 481-493. <https://www.ncbi.nlm.nih.gov/pubmed/17381374>

Studies Discussed During the Episode

MacLaren R, et al. Effects of pharmacist participation in intensive care units on clinical and economic outcomes of critically ill patients with thromboembolic or infarction-related events. *Pharmacotherapy* 2009; 29(7): 761-768. <https://www.ncbi.nlm.nih.gov/pubmed/19558249>

Rivkin A, et al. Evaluation of the role of the critical care pharmacist in identifying and avoiding or minimizing significant drug-drug interactions in medical intensive care patients. *J Crit Care* 2011; 26(1): e1-e6. <https://www.ncbi.nlm.nih.gov/pubmed/20646898>

Leape LL, et al. Pharmacist participation on physician rounds and adverse drug events in the intensive care unit. *JAMA* 1999; 282(3): 267-270. <https://www.ncbi.nlm.nih.gov/pubmed/10422996>

MacLaren R, et al. Clinical and economic outcomes of involving pharmacists in the direct care of critically ill patients with infections. *Crit Care Med* 2008; 36(12): 3184-3189. <https://www.ncbi.nlm.nih.gov/pubmed/18936700>

Marshall J, et al. Impact of a clinical pharmacist-enforced intensive care unit sedation protocol on duration of mechanical ventilation and hospital stay. *Crit Care Med* 2008; 36(2): 427-433. <https://www.ncbi.nlm.nih.gov/pubmed/18091554>

Louzon P, et al. Impact of pharmacist management of pain, agitation, and delirium in the intensive care unit through participation in multidisciplinary bundle rounds. *Am J Health Syst Pharm* 2017; 74(4): 253-262. <https://www.ncbi.nlm.nih.gov/pubmed/28179250>

Hahn L, et al. Pharmacist-developed sedation protocol and impact on ventilator days. *J Pharm Pract* 2013; 26(4): 406-408. <https://www.ncbi.nlm.nih.gov/pubmed/23204145>

Nguyen CT, et al. Pharmacists' Impact on Glycemic Control Among Surgical Patients at a Large Academic Institution. *Clin Diabetes* 2016; 34(2): 105-108.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4833484/>

Bond CA, et al. Pharmacist-provided anticoagulation management in United States hospitals: death rates, length of stay, Medicare charges, bleeding complications, and transfusions. *Pharmacotherapy* 2004; 24: 953-963. <https://www.ncbi.nlm.nih.gov/pubmed/15338843>

Flynn JD, et al. Utilization of Pharmacist Responders as a Component of a Multidisciplinary Sepsis Bundle. *Ann Pharmacother* 2014; 48(9): 1145-1151.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4376659/>

Buckley MS, et al. Impact of a clinical pharmacist stress ulcer prophylaxis management program on inappropriate use in hospitalized patients. *Am J Med* 2015; 128(8): 905-913.
<https://www.ncbi.nlm.nih.gov/pubmed/25820164>

Oxman DA, et al. Improving antibiotic de-escalation in suspected ventilator-associated pneumonia: an observational study with a pharmacist-driven intervention. *J Pharm Pract* 2015; 28(5): 457-461. <https://www.ncbi.nlm.nih.gov/pubmed/24651641>

Lingenfelter E, et al. ED pharmacist monitoring of provider antibiotic selection aids appropriate treatment for outpatient UTI. *Am J Emerg Med* 2016; 34(8): 1600-1603.
<https://www.ncbi.nlm.nih.gov/pubmed/27306262>

Rech MA, et al. Pharmacist Participation in Acute Ischemic Stroke Decreases Door-To-Needle Time to Recombinant Tissue Plasminogen Activator. *Ann Pharmacother* 2017; 51(12): 1084-1089. <https://www.ncbi.nlm.nih.gov/pubmed/28759998>

Patanwala AE, et al. Pharmacist's activities on a trauma response team in the emergency department. *Am J Health Syst Pharm* 2010; 67(18): 1536-1538.
<https://www.ncbi.nlm.nih.gov/pubmed/20811031>

Robey-Gavin E. Impact of Clinical Pharmacists on Initiation of Postintubation Analgesia in the Emergency Department. *J Emerg Med* 2016; 50(2): 308-314.
<https://www.ncbi.nlm.nih.gov/pubmed/26433427>

Amini A, et al. Effect of a pharmacist on timing of postintubation sedative and analgesic use in trauma resuscitations. *Am J Health Syst Pharm* 2013; 70(17): 1513-1517.

<https://www.ncbi.nlm.nih.gov/pubmed/23943183>

Masic D, et al. Pharmacist Presence Decreases Time to Prothrombin Complex Concentrate in Emergency Department Patients with Life-Threatening Bleeding and Urgent Procedures. *J Emerg Med* 2019; pii: S0736-4679(19)30488-3. doi: 10.1016/j.jemermed.2019.06.027. [Epub ahead of print]. <https://www.ncbi.nlm.nih.gov/pubmed/31447188>

Hammond DA, et al. Scoping review of interventions associated with cost avoidance able to be performed in the Intensive Care Unit and Emergency Department. *Pharmacotherapy* 2019; 39(3): 215-231. <https://www.ncbi.nlm.nih.gov/pubmed/30664269>

Hammond DA, et al. Cost avoidance associated with clinical pharmacist presence in a medical intensive care unit. *J Am Coll Clin Pharm* 2019; **(**): 1-6.
<https://accpjournals.onlinelibrary.wiley.com/doi/10.1002/jac5.1111>

Dasta JF. Critical Care. *Ann Pharmacother* 2006; 40(4): 736-737.
<https://www.ncbi.nlm.nih.gov/pubmed/16569809>