

Vasoactives in Septic Shock References

Special Guest: Alex Flannery, PharmD, BCCCP, BCPS

Guidelines/General Reviews

Singer M., et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016; 315(8): 801-810. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4968574/>

Rhodes A., et al. Surviving Sepsis Campaign: International Guidelines for the Management of Sepsis and Septic Shock: 2016. *Intensive Care Med* 2017; 43(3): 304-377. <https://www.ncbi.nlm.nih.gov/pubmed/28101605>

Levy M.M., et al. The Surviving Sepsis Campaign Bundle: 2018 Update. *Crit Care Med* 2018; 46(6): 997-1000. <https://www.ncbi.nlm.nih.gov/pubmed/29767636>

Angus D.C., et al. Severe sepsis and septic shock. *N Eng J Med* 2013; 369(9): 840-851. <https://www.ncbi.nlm.nih.gov/pubmed/23984731>

Dalimonte M.A., et al. Vasoactive Agents for Adult Septic Shock: An Update and Review. *J Pharm Pract* 2019; 897190019844124. <https://www.ncbi.nlm.nih.gov/pubmed/31057085>

Hollenberg S.M. Vasoactive drugs in circulatory shock. *Am J Respir Crit Care Med* 2011; 183(7): 847-855. <https://www.ncbi.nlm.nih.gov/pubmed/21097695>

Studies Discussed During the Episode

Polito A., et al. Pharmacokinetics of oral fludricortisone in septic shock. *Br J Clin Pharmacol* 2016; 82(6): 1509-1516. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5099539/>

Annane D. Why My Steroid Trials Were “Positive”? *Crit Care Med* 2019; doi: 10.1097/CCM.0000000000003889. [Epub ahead of print] <https://www.ncbi.nlm.nih.gov/pubmed/31259754>

Marik P.E., et al. Hydrocortisone, Vitamin C, and Thiamine for the Treatment of Severe Sepsis and Septic Shock: A Retrospective Before-After Study. *Chest* 2017; 151(6): 1229-1238. <https://www.ncbi.nlm.nih.gov/pubmed/27940189>

Donnino M.W., et al. Randomized, Double-Blind, Placebo-Controlled Trial of Thiamine as a Metabolic Resuscitator in Septic Shock: A Pilot Study. *Crit Care Med* 2016; 44(2): 360-367. <https://www.ncbi.nlm.nih.gov/pubmed/26771781>

Woolum J.A., et al. Effect of Thiamine Administration on Lactate Clearance and Mortality in Patients with Septic Shock. *Crit Care Med* 2018; 46(11): 1747-1752. <https://www.ncbi.nlm.nih.gov/pubmed/30028362>

Khanna A., et al. Angiotensin II for the Treatment of Vasodilatory Shock. *N Eng J Med* 2017; 377(5): 419-430. <https://www.ncbi.nlm.nih.gov/pubmed/28528561>

Wong A., et al. A Retrospective Review of Angiotensin II Use in Adult Patients with Refractory Distributive Shock. *J Intensive Care Med* 2019; 885066619872720. doi: 10.1177/0885066619872720. [Epub ahead of print]. <https://www.ncbi.nlm.nih.gov/pubmed/31480886>

Bissell B.D., et al. A Blast From the Past: Revival of Angiotensin II for Vasodilatory Shock. *Ann Pharmacother* 2018; 52(9): 920-927. <https://www.ncbi.nlm.nih.gov/pubmed/29582666>

Paciullo C.A., et al. Methylene blue for the treatment of septic shock. *Pharmacotherapy* 2010; 30(7): 702-715. <https://www.ncbi.nlm.nih.gov/pubmed/20575634>

Vail E., et al. Association Between US Norepinephrine Shortage and Mortality Among Patients with Septic Shock. *JAMA* 2017; 317(14): 1433-1442. <https://www.ncbi.nlm.nih.gov/pubmed/28322415>

Hammond D.A., et al. Effects of Norepinephrine and Vasopressin Discontinuation Order in the Recovery Phase of Septic Shock: A Systematic Review and Individual Patient Data Meta-Analysis. *Pharmacotherapy* 2019; 39(5): 544-552. <https://www.ncbi.nlm.nih.gov/pubmed/30893494>

Rizvi M.S., et al. Trends in Use of Midodrine in the ICU: A Single-Center Retrospective Case Series. *Crit Care Med* 2018; 46(7): e628-e633. <https://www.ncbi.nlm.nih.gov/pubmed/29613861>

Future and In-Progress Research

Crystalloid Liberal or Vasopressors Early Resuscitation in Sepsis (CLOVERS) trial. <https://clinicaltrials.gov/ct2/show/NCT03434028>

Coopersmith C.M., et al. Research Priorities for Sepsis and Septic Shock. *Crit Care Med* 2018; 46(8): 1334-1356. <https://www.ncbi.nlm.nih.gov/pubmed/29957716>