

Mar/Apr 2021 Literature Review References

Special Guests: Ashley Barlow, PharmD & Brooke Barlow, PharmD

6-Pack of Studies

Roquilly A, Moyer JD, Huet O, et al. Effect of continuous infusion of hypertonic saline vs standard care on 6-month neurological outcomes in patients with traumatic brain injury: the COBI randomized clinical trial. *JAMA*. 2021;325(20):2056.

<https://pubmed.ncbi.nlm.nih.gov/34032829/>

Khan AA, Gurnani PK, Peksa GD, Whittier WL, DeMott JM. Bisphosphonate versus bisphosphonate and calcitonin for the treatment of moderate to severe hypercalcemia of malignancy. *Ann Pharmacother*. 2021;55(3):277-285.

<https://pubmed.ncbi.nlm.nih.gov/32885992/>

Russotto V, Myatra SN, Laffey JG, et al. Intubation practices and adverse peri-intubation events in critically ill patients from 29 countries. *JAMA*. 2021;325(12):1164.

<https://pubmed.ncbi.nlm.nih.gov/33755076/>

Burns KEA, Rizvi L, Cook DJ, et al. Ventilator weaning and discontinuation practices for critically ill patients. *JAMA*. 2021; 325(12): 1173-1184.

<https://pubmed.ncbi.nlm.nih.gov/33755077/>

Bosch NA, Rucci JM, Massaro JM, et al. Comparative effectiveness of heart rate control medications for the treatment of sepsis-associated atrial fibrillation. *Chest*. 2021;159(4):1452-1459. <https://pubmed.ncbi.nlm.nih.gov/33619010/>

Rivosecchi RM, Arakelians AR, Ryan J, et al. Comparison of anticoagulation strategies in patients requiring venovenous extracorporeal membrane oxygenation: heparin versus bivalirudin. *Critical Care Medicine*. 2021; 49(7): 1129-1136. <https://pubmed.ncbi.nlm.nih.gov/33711003/>

Seelhammer TG, Bohman JK, Schulte PJ, Hanson AC, Aganga DO. Comparison of bivalirudin versus heparin for maintenance systemic anticoagulation during adult and pediatric extracorporeal membrane oxygenation. *Critical Care Medicine*. 2021 Apr 19. doi: 10.1097/CCM.0000000000005033. Online ahead of print.

<https://pubmed.ncbi.nlm.nih.gov/33870916/>

Fisser C, Winkler M, Malfertheiner MV, et al. Argatroban versus heparin in patients without heparin-induced thrombocytopenia during venovenous extracorporeal membrane oxygenation: a propensity-score matched study. *Crit Care*. 2021;25(1):160.

<https://pubmed.ncbi.nlm.nih.gov/33910609/>

China L, Freemantle N, Forrest E, et al. A randomized trial of albumin infusions in hospitalized patients with cirrhosis. *N Engl J Med*. 2021;384(9):808-817.

<https://pubmed.ncbi.nlm.nih.gov/33657293/>

Sepsis Studies

Taylor SP, Anderson WE, Beam K, Taylor B, Ellerman J, Kowalkowski MA. The association between antibiotic delay intervals and hospital mortality among patients treated in the emergency department for suspected sepsis. *Critical Care Medicine*. 2021;49(5):741-747.

<https://pubmed.ncbi.nlm.nih.gov/33591002/>

Demiselle J, Seegers V, Lemerle M, et al. Prior exposure to angiotensin ii receptor blockers in patients with septic shock to individualize mean arterial pressure target? A post hoc analysis of the sepsis and mean arterial pressure (Sepsispam) trial. *Critical Care Medicine*.

2021;49(4):e412-e422. <https://pubmed.ncbi.nlm.nih.gov/33591012/>

Jörres A. Initial arterial blood pressure targets in patients with septic shock: one size fits all or made to measure? *Critical Care Medicine*. 2021;49(4):719-721.

<https://pubmed.ncbi.nlm.nih.gov/33731615/>

Lanspa MJ, Cirulis MM, Wiley BM, et al. Right ventricular dysfunction in early sepsis and septic shock. *Chest*. 2021;159(3):1055-1063. <https://pubmed.ncbi.nlm.nih.gov/33068615/>

Jagan N, Morrow LE, Walters RW, et al. Sepsis, the administration of iv fluids, and respiratory failure. *Chest*. 2021;159(4):1437-1444. <https://pubmed.ncbi.nlm.nih.gov/33197405/>

Lat I, Coopersmith CM, De Backer D. The surviving sepsis campaign: fluid resuscitation and vasopressor therapy research priorities in adult patients. *Critical Care Medicine*. 2021;49(4):623-635. <https://pubmed.ncbi.nlm.nih.gov/33731607/>

Pain, Agitation, Delirium, and Neuromuscular Blockade

Duprey MS, Devlin JW, van der Hoeven JG, et al. Association between incident delirium treatment with haloperidol and mortality in critically ill adults. *Critical Care Medicine*. 2021; 49(8): 1303-1311. <https://pubmed.ncbi.nlm.nih.gov/33861548/>

Pappal RD, Roberts BW, Winkler W, Yaegar LH, Stephens RJ, Fuller BM. Awareness with paralysis in mechanically ventilated patients in the emergency department and icu: a systematic review and meta-analysis*. *Critical Care Medicine*. 2021;49(3):e304-e314. <https://pubmed.ncbi.nlm.nih.gov/33566462/>

Wongtangman K, Grabitz SD, Hammer M, et al. Optimal sedation in patients who receive neuromuscular blocking agent infusions for treatment of acute respiratory distress syndrome—a retrospective cohort study from a new england health care network. *Critical Care Medicine*. 2021; 49(7): 1137-1148. <https://pubmed.ncbi.nlm.nih.gov/33710031/>

Romero N, Dube KM, Lupi KE, DeGrado JR. Evaluation of delirium in critically ill patients prescribed melatonin or ramelteon. *Ann Pharmacother*. 2021 Mar 13; 10600280211002054. doi: 10.1177/10600280211002054. Online ahead of print. <https://pubmed.ncbi.nlm.nih.gov/33715466/>

Shehabi Y, Serpa Neto A, et al. Early sedation with dexmedetomidine in ventilated critically ill patients and heterogeneity of treatment effect in the SPICE III randomised controlled trial. *Intensive Care Med*. 2021;47(4):455-466. <https://pubmed.ncbi.nlm.nih.gov/33686482/>

Park JH, Derry KL, Owens RL. Dexmedetomidine as add-on sedation to reduce continuous infusion sedative use in mechanically ventilated patients. *American Journal of Health-System Pharmacy*. 2021;78(Supplement_1):S1-S9. <https://pubmed.ncbi.nlm.nih.gov/33038212/>

von Oelreich E, Eriksson M, Sjölund K-F, Discacciati A, Larsson E, Oldner A. Opioid use after intensive care: a nationwide cohort study. *Critical Care Medicine*. 2021;49(3):462-471. <https://pubmed.ncbi.nlm.nih.gov/33512940/>

Nazarnia S, Subramaniam K. Nonopioid analgesics in postoperative pain management after cardiac surgery. *Semin Cardiothorac Vasc Anesth*. 2021; 1089253221998552. doi: 10.1177/1089253221998552. Online ahead of print. <https://pubmed.ncbi.nlm.nih.gov/33899581/>

Cicci CD, Fudzie SS, Campbell-Bright S, et al. Accuracy and safety of medication histories obtained at the time of intensive care unit admission of delirious or mechanically ventilated patients. *Am J Health-Syst Pharm*. 2021; 78(8): 736-742. <https://pubmed.ncbi.nlm.nih.gov/33580666/>

Gélinas C, Bérubé M, Puntillo KA, et al. Validation of the Critical-Care Pain Observation Tool-Neuro in brain-injured adults in the intensive care unit: a prospective cohort study. *Crit Care*. 2021;25(1):142. <https://pubmed.ncbi.nlm.nih.gov/33849619/>

Infectious Diseases and Pharmacokinetic Studies

Narayan SW, Thoma Y, Drennan PG, et al. Predictive performance of bayesian vancomycin monitoring in the critically ill . *Crit Care Med*. 2021 Apr 28. doi: 10.1097/CCM.0000000000005062. Online ahead of print. <https://pubmed.ncbi.nlm.nih.gov/33938713/>

Alvarez-Arango S, Ogunwale SM, Sequist TD, Burk CM, Blumenthal KG. Vancomycin infusion reaction — moving beyond “red man syndrome.” *N Engl J Med*. 2021;384(14):1283-1286. <https://pubmed.ncbi.nlm.nih.gov/33830710/>

The Azurea Network, Roger C, Louart B, et al. An international survey on aminoglycoside practices in critically ill patients: the AMINO III study. *Ann Intensive Care*. 2021;11(1):49. <https://pubmed.ncbi.nlm.nih.gov/33740157/>

Al-Shaer MH, Philpott CD, Droege CA, et al. Cefepime population pharmacokinetics and target attainment in critically ill patients on continuous renal replacement therapy. *Antimicrob Agents Chemother*. 2021; 65(6): e00144-21. <https://pubmed.ncbi.nlm.nih.gov/33722885/>

El-Haffaf I, Caissy J-A, Marsot A. Piperacillin-tazobactam in intensive care units: a review of population pharmacokinetic analyses. *Clin Pharmacokinet*. 2021; 60(7): 855-875. <https://pubmed.ncbi.nlm.nih.gov/33876381/>

Roberts JA, Joynt GM, Lee A, et al. The effect of renal replacement therapy and antibiotic dose on antibiotic concentrations in critically ill patients: data from the multinational sampling antibiotics in renal replacement therapy study. *Clinical Infectious Diseases*. 2021;72(8):1369-1378. <https://pubmed.ncbi.nlm.nih.gov/32150603/>

Howatt M, Klompas M, Kalil AC, Metersky ML, Muscedere J. Carbapenem antibiotics for the empiric treatment of nosocomial pneumonia. *Chest*. 2021;159(3):1041-1054. <https://pubmed.ncbi.nlm.nih.gov/33393468/>

Hites M. Minireview on novel anti-infectious treatment options and optimized drug regimens for sepsis. *Front Med*. 2021; 8: 640740. <https://pubmed.ncbi.nlm.nih.gov/33937283/>

Sousa-Pinto B, Blumenthal KG, Courtney L, Mancini CM, Jeffres MN. Assessment of the frequency of dual allergy to penicillins and cefazolin: a systematic review and meta-analysis. *JAMA Surg.* 2021;156(4):e210021. <https://pubmed.ncbi.nlm.nih.gov/33729459/>

Tamma PD, Aitken SL, Bonomo RA, Mathers AJ, van Duin D, Clancy CJ. Infectious diseases society of america guidance on the treatment of extended-spectrum β -lactamase producing enterobacterales (Esbl-e), carbapenem-resistant enterobacterales (Cre), and pseudomonas aeruginosa with difficult-to-treat resistance (DTR- P. aeruginosa). *Clinical Infectious Diseases.* 2021;72(7):1109-1116. <https://pubmed.ncbi.nlm.nih.gov/33830222/>

Front of the Fridge (Pharmacist-Driven Studies)

Erstad BL. Usefulness of the biomarker TIMP-2•IGFBP7 for acute kidney injury assessment in critically ill patients: a narrative review. *Ann Pharmacother.* Published online April 8, 2021:106002802110054. <https://pubmed.ncbi.nlm.nih.gov/33829897/>

Dubrawka CA, Betthauser KD, Pope HE, Gibson GA. Effect of vasopressin dose on hemodynamic response in obese patients with septic shock: a retrospective observational study. *Ann Pharmacother.* Published online April 3, 2021:106002802110072. <https://pubmed.ncbi.nlm.nih.gov/33813908/>

Webb AJ, Brown CS, Naylor RM, Rabinstein AA, Mara KC, Nei AM. Thromboelastography is a marker for clinically significant progressive hemorrhagic injury in severe traumatic brain injury. *Neurocrit Care.* 2021 Apr 12. Doi: 10.1007/s12028-021-01217-0. Online ahead of print. <https://pubmed.ncbi.nlm.nih.gov/33846901/>

Buckley MS, Agarwal SK, Garcia-Orr R, Saggat R, MacLaren R. Comparison of fixed-dose inhaled epoprostenol and inhaled nitric oxide for acute respiratory distress syndrome in critically ill adults. *J Intensive Care Med.* 2021;36(4):466-476. <https://pubmed.ncbi.nlm.nih.gov/32133901/>

Cascone AE, Daley MJ, Pan N, Padilla-Tolentino E, Milling TJ. Low dose versus standard dose four factor prothrombin complex concentrate for factor-Xa inhibitor reversal in spontaneous and traumatic intracranial hemorrhage. *Pharmacotherapy.* 2021; 41(6): 501-507. <https://pubmed.ncbi.nlm.nih.gov/33866591/>

Hunt AR, Coffeen SN, Shiltz DL, Ice C, Parker J. Factor viii inhibitor bypassing activity (Feiba) reversal for apixaban and rivaroxaban in patients with acute intracranial and nonintracranial hemorrhage. *Ann Pharmacother.* Published online April 10, 2021:106002802110045. <https://pubmed.ncbi.nlm.nih.gov/33843267/>

Bennett S, Mullen C, Mistry B, Cucci M. Performance of nasal methicillin-resistant *Staphylococcus aureus* screening for intra-abdominal infections in critically ill adult patients. *Pharmacotherapy*. 2021;41(3):257-264. <https://pubmed.ncbi.nlm.nih.gov/33345308/>

Murphy JA, Curran BM, Gibbons WA, Harnica HM. Adjunctive phenobarbital for alcohol withdrawal syndrome: a focused literature review. *Ann Pharmacother*. Published online March 7, 2021:1060028021999821. <https://pubmed.ncbi.nlm.nih.gov/33678057/>

Erstad BL. Predicted body weight: A rose by any other name. *American Journal of Health-System Pharmacy*. 2021;78(8):751-753. <https://pubmed.ncbi.nlm.nih.gov/33580244/>

Kane-Gill SL, Kaplan LJ. Key roles for pharmacists in family-centered care to advance patient safety. *Critical Care Medicine*. 2021;49(4):e454-e455. <https://pubmed.ncbi.nlm.nih.gov/33731616/>