

Trial of the Week: CATS

Special Guest: Scott Benken, PharmD, MHPE,
BCPS-AQ Cardiology, FCCM, FCCP

CATS trial

Annane D, Vignon P, Renault A, Bollaert PE, Charpentier C, Martin C, Troché G, Ricard JD, Nitenberg G, Papazian L, Azoulay E, Bellissant E; CATS Study Group. Norepinephrine plus dobutamine versus epinephrine alone for management of septic shock: a randomised trial. *Lancet*. 2007 Aug 25;370(9588):676-84. <https://pubmed.ncbi.nlm.nih.gov/17720019/>

Articles/Guidelines Referenced

Hollenberg SM, Ahrens TS, Annane D, Astiz ME, Chalfin DB, Dasta JF, Heard SO, Martin C, Napolitano LM, Susla GM, Totaro R, Vincent JL, Zanotti-Cavazzoni S. Practice parameters for hemodynamic support of sepsis in adult patients: 2004 update. *Crit Care Med*. 2004 Sep;32(9):1928-48. <https://pubmed.ncbi.nlm.nih.gov/15343024/>

Annane D, Sébille V, Charpentier C, et al. Effect of treatment with low doses of hydrocortisone and fludrocortisone on mortality in patients with septic shock. *JAMA*. 2002 Aug 21;288(7):862-71. <https://pubmed.ncbi.nlm.nih.gov/12186604/>

Evans L, Rhodes A, Alhazzani W, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock 2021. *Crit Care Med*. 2021 Nov 1;49(11):e1063-e1143. <https://pubmed.ncbi.nlm.nih.gov/34605781/>